



**PERMIT TO ALLOW MINOR USE OF A REGISTERED AGVET CHEMICAL PRODUCT
FOR CONTROL OF CERCOSPORA LEAF SPOT DISEASE IN PERSIMMONS**

PERMIT NUMBER – PER12488

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows any person to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 27 OCTOBER 2010 TO 31 MARCH 2025

Permit Holder:

HORTICULTURE INNOVATION AUSTRALIA LTD
Level 7, 141 Walker Street
NORTH SYDNEY NSW 2060

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Product to be used:

FARMOZ MANCOZEB 800 AGRICULTURAL FUNGICIDE (APVMA No. 39325)

PLUS OTHER REGISTERED PRODUCTS

Containing: 800 g/kg MANCOZEB as the only active constituent.

SABERO MANCOZEB 750DF FUNGICIDE (APVMA No. 54451)

PLUS OTHER REGISTERED PRODUCTS

Containing: 750 g/kg MANCOZEB as the only active constituent.

Directions for Use:

Crop Type	Plant Disease	Application Rate
PERSIMMON	CERCOSPORA OR ANGULAR LEAF SPOT (<i>Cercospora kaki</i>)	<u>Dilute Spray:</u> 200 g product/100 L <u>Concentrate spray:</u> Up to five times (5x) Refer to Mixing/Application Section.

Critical Use Comments:

- DO NOT apply more than 14 times in one season.
- DO NOT apply more than 6 applications after shuck fall.
- Cover spray every 2-4 weeks from bud-break to harvest at intervals of not less than 14 days.
- Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.

MIXING/APPLICATION:

Dilute spraying:

- Apply by dilute spraying equipment. Use a sprayer designed to apply high volumes of water up to the point of runoff and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different setting on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate spraying:

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

Example only

- Dilute spray volume as determined above: For example 1500 L/ha
- Your chosen concentrate spray volume: For example 500 L/ha

- The concentration factor in this example is: 3 X (i.e. $1500\text{ L} \div 500\text{ L} = 3$)
- If the dilute label rate is 10 mL/100 L, then the concentrate rate becomes 3 X 10, that is 30 mL/100 L of concentrate spray.
- The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
- Do not use a concentrate rate higher than that specified in the Critical Comments.
- For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry best practices.

Withholding Period:

DO NOT harvest for 14 DAYS after application.

Jurisdiction:

All States and Territories, except VIC.

Note: Victoria is not included in this permit, as their Control-of-Use legislation means a permit is not required to legalise this off-label use in that State.

Additional Conditions:

This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the product label.

Export of treated produce

MRLs can be found in the *Agricultural and Veterinary Chemicals Code (MRL Standard) Instrument 2019*. MRLs apply only to produce marketed and consumed in Australia. If treated produce is to be exported, residues must not exceed the limits/tolerances of the importing country.

Issued by the Australian Pesticides and Veterinary Medicines Authority

Note: 28/03/2020 – Permit holder updated. Expiry extended to 31 March 2025. Permit issued as Version 3.